

Evoked thrill: A simple intraoperative maneuver predicts early patency of arteriovenous fistulas

Miltos K. Lazarides, MD, Demetrios N. Staramos, MD, Vasilios D. Tzilalis, MD, Konstantinos E. Simopoulos, MD, and John N. Dayantas, MD, *Athens and Alexandroupolis, Greece*

Arteriovenous fistula at the wrist has remained the procedure of choice for long-term chronic hemodialysis since its development by Brescia and Cimino in 1966.¹ However, the early failure rate (<30 days) remains a significant problem, affecting 24% of patients in a series reported by Palder.² Beyond the first month, the long-term patency rate for radiocephalic fistulas is excellent.

Compromised venous outflow from the fistula, usually because of preexisting areas of segmental stenotic lesions, is the most common cause of early failure. The cephalic vein must be investigated all the way from the wrist by clinical inspection, preoperative color Doppler imaging, intraoperative venography, or angiography.^{3,4} We describe a simple intraoperative maneuver that can be used to detect proximal stenotic vein segments.

METHODS

The radial artery and cephalic vein are exposed and mobilized through a standard longitudinal incision parallel with the lateral border of the forearm. A venotomy approximately 0.8 cm long is performed, and a small plastic tube (e.g., 14-gauge catheter with a 2 mm external diameter) is gently inserted through the opening. The catheter is connected to a syringe containing heparinized physiologic saline, and the surgeon injects 20 ml of saline while holding the syringe with his left hand. The saline is injected by applying pressure intermittently, mimicking the

pulsed arterial flow. While injecting saline, the surgeon palpates the skin over the proximal vein with his right hand, and if there is no stenotic lesion, an evoked thrill is felt (Fig. 1). When a palpable thrill is produced, the surgeon proceeds with an arteriotomy in the radial artery and performs an arteriovenous anastomosis. When no thrill is produced, the surgeon should suspect the existence of a stenotic lesion or areas of thickened vein wall. For these cases, "on table" venography is performed to reveal the exact nature of the lesion, and unnecessary arteriotomy of the radial artery is avoided. After the saline injection, the tube is left in place to avoid occluding the vein with vascular clamps. The proximal vein is irrigated periodically with heparinized saline solution through the catheter threaded up its lumen, ensuring patency while a vessel loop placed around the intubated vein prevents backbleeding. The distal venous limb is always ligated on completion of the anastomosis to avoid venous hypertension of the hand.

RESULTS

From November 1996, when the evoked thrill maneuver was introduced as a routine technique in every autologous arteriovenous fistula procedure (except basilic transpositions), through August 1997, 49 consecutive patients who underwent 52 autologous access operations were identified by retrospective review of their records. The clinical details of the patients in the study group are summarized in Table I.

The evoked thrill maneuver was performed in all 49 patients. When the thrill was produced, the test result was considered negative (i.e., proximal vein stenosis was unlikely). In the absence of an induced palpable thrill, the test result was considered positive (i.e., proximal vein stenosis was likely). The early failure rate for the series was 17%, with nine arteriovenous fistulas thrombosed within the first postoperative month. For four of the nine early failures, the evoked thrill test results were positive. The

From the Department of Vascular Surgery (Drs. Lazarides, Staramos, Tzilalis, and Dayantas), Athens General Hospital "G Gennimatas" and from the Second Department of Surgery (Drs. Staramos and Simopoulos), Thrace Democritus University, Alexandroupolis.

Reprint requests: Miltos K. Lazarides, Amaseias 5, Athens 11634, Greece.

J Vasc Surg 1998;27:750-2.

Copyright © 1998 by the Society for Vascular Surgery and International Society for Cardiovascular Surgery, North American Chapter.

0741-5214/98/\$5.00 + 0 24/4/87095




Fig. 1. An evoked thrill is palpable with the surgeon's right hand while he is injecting the saline through an intravenous catheter and holding the syringe with his left hand. The saline is injected by applying pressure intermittently, mimicking the pulsed arterial flow.

Table I. Clinical details of 49 consecutive patients who underwent 52 autologous arteriovenous fistula procedures

Age (yr)	mean = 57.5; range, 16–83
Male/female	32/17
Access type	
Radiocephalic fistulas at wrist	31
Radiocephalic fistulas at forearm	10
Brachiocephalic fistulas at elbow	11
Primary/secondary procedures*	42/10

*A procedure was considered secondary when a previous access had failed in the ipsilateral limb.

decision to proceed with the procedure in these patients despite a positive test result was based on free and easy passage of the catheter through the vein from the wrist to a level above the antecubital fossa. The existence of a proximal stenotic, nondistensible vein segment with thickened intimal surface was revealed by direct dissection when the access was revised in two patients and in a third patient by table venography. In the latter case, no revision was performed. The cause of early failure was not disclosed for all patients with negative test results, but in two it correlated with premature puncture during the second postoperative week. Table II compares the results of the evoked thrill maneuver with the early failure rate.

No false-positive test results occurred. The early failure rate is a better criterion of efficacy of the maneuver than long-term patency. The latter is affected by factors such as puncture timing and technique and by development of neointimal hyperplasia.

Table II. Comparison of evoked thrill maneuver results with the early failure rate for 52 autologous arteriovenous fistulas

Early failure (<30 d)	Evoked thrill maneuver*	
	Negative responses	Positive responses
Yes	5	4
No	43	0

*For the evoked thrill maneuver: sensitivity = 44%, specificity = 100%, positive predictive value = 100%, and negative predictive value = 89%.

DISCUSSION

Thrombosis is the most common reason for early arteriovenous fistula loss. Thromboses within the first 4 to 6 weeks postoperatively are caused by technical problems at the anastomosis or stenotic lesions of the more proximal vein.⁵ The latter are caused by prior episodes of phlebitis and usually are not recognized preoperatively. The most common lesions occur in regions of previous thrombosis and recanalization; these are webs, strands, or areas of nondistensible vein with a thickened intimal surface.

Proximal stenotic lesions should be recognized preoperatively or intraoperatively, but because no consensus exists regarding the ideal preoperative screening method for access surgery, most surgeons are satisfied with preoperative clinical evaluation of the vein, including tourniquet application and vein percussion.⁵ However, preoperative clinical examination, with or without a tourniquet, and direct intraoperative inspection of the vein frequently fail to detect proximal stenotic lesions. Preoperative Duplex scanning or intraoperative angiography are more accurate methods for assessing the quality of venous outflow. Duplex scanning has an 88% specificity for detecting the suitability of the arm veins for access surgery,³ but not all centers performing vascular access are equipped with color Doppler or angiography. Because angiography requires clearing the vein of blood, some irrigation fluid is necessary, but some patients cannot tolerate this procedure.⁴

Presence of a palpable thrill postoperatively is the best indicator of good venous runoff, whereas the absence of a thrill and the existence of strong pulsations of the proximal vein suggest outflow obstruction and demand access revision.⁶ The evoked thrill is a simple method that can be used to reproduce flow conditions and prevent inadvertent arteriotomy and subsequent compromise of distal perfusion. In

our series the evoked thrill maneuver, with a sensitivity of only 44%, failed to locate about one half of the early failures. The reason appears to be that early failure is not always caused by a proximal vein stenotic lesion; systemic pressure drop, premature cannulation, and inadequate inflow also cause early failures. However, the positive predictive value (of an early failure) of the test was 100%, indicating that this maneuver is highly accurate in predicting early thrombosis and constitutes a simple and useful intra-operative method for improving the patency rate of autologous arteriovenous fistulas.

REFERENCES

1. Brescia MJ, Cimino JE, Appel H, Hurwich BJ. Chronic hemodialysis using venipuncture and a surgically created arteriovenous fistula. *N Engl J Med* 1966;275:1089-92.
2. Palder SB, Kirkman RL, Whittemore AD, Hakim RM, Lazarus JM, Tilney NL. Vascular access for hemodialysis, patency rates and results of revision. *Ann Surg* 1985;202:235-9.
3. Tordoir JHM. Duplex scanning of hemodialysis access operations. In: Bernstein EF, editor. *Vascular diagnosis*. 4th ed. St. Louis, Mosby; 1993:599-607.
4. Miller A, Holzenbein TJ, Gottlieb MN. Routine angioscopy for vascular access surgery. In: Henry ML, Ferguson RM, editors. *Vascular access for hemodialysis-IV*. Precept Press; 1995:94-102.
5. Ryan JJ, Dennis MJS. Radiocephalic fistula in vascular access. *Br J Surg* 1990;77:1321-2.
6. Marx AB, Landmann J, Harder FH. Surgery for vascular access. *Curr Probl Surg* 1990;27:5-48.

Submitted July 9, 1997; accepted Oct. 23, 1997.

Announcing a Home Page on the WWW for the Vascular Surgical Societies

To enter the exciting new world of cyberspace, simply point your computer to :

<http://www.vascsurg.org>

and hang on to your hat! You can scan back issues of the *Journal of Vascular Surgery*, look up a colleague who is a member of most regional vascular societies, review abstracts for upcoming vascular meetings, analyze a challenging "Case of the Month," and enjoy many other interesting features.

Don't forget to visit the "Welcome" area for the latest information on navigating the site, and please register for your user name and password if you have not already received these as a member of either The Society for Vascular Surgery or the North American Chapter of the International Society for Cardiovascular Surgery.

SEE YOU ON THE WEB!

Richard F. Kempczinski, MD
WebMaster